

● **TOYOBO ENZYMES** ●
(Diagnostic Reagent Grade)

D-LACTATE DEHYDROGENASE

from Microorganism

(R)-Lactate:NAD⁺ oxidoreductase (EC 1. 1. 1. 28)



PREPARATION and SPECIFICATION

Appearance	: White amorphous powder, lyophilized	
Activity	: Grade II 900U/mg-solid or more	
Contaminants	: NADH oxidase	≤1.0×10 ⁻³ %
	Malate dehydrogenase	≤1.0×10 ⁻² %
	GOT	≤5.0×10 ⁻³ %
	GPT	≤5.0×10 ⁻³ %
	Myokinase	≤1.0×10 ⁻² %
	Pyruvate kinase	≤1.0×10 ⁻³ %

PROPERTIES

Stability	: Stable at -20°C	
Molecular weight	: approx. 140,000	
Isoelectric point	: 4.0	
Michaelis constants	: 6.4×10 ⁻⁴ M (pyruvate, pH 7.0)	
Inhibitors	: Ag ²⁺ , Hg ²⁺ , SH-reagents	
Optimum pH	: 5.0—7.0	(Fig.2)
Optimum temperature	: 30—37°C	(Fig.3)
pH Stability	: pH 5.0—9.0 (25°C, 48hr)	(Fig.4)
Thermal stability	: below 45°C (pH 7.0, 15min)	(Fig.5)
Effect of various chemicals	: (Table 1)	

APPLICATIONS

This enzyme is useful for enzymatic determination of numerous metabolites, e.g. ATP, ADP, glucose, creatinine, pyruvate, lactate and glycerol, and of enzyme activities, e.g. GPT, PK, and CPK when coupled with the related enzymes.

ASSAY

Principle:



The disappearance of NADH is measured at 340nm by spectrophotometry.

Unit definition:

One unit causes the oxidation of one micromole of NADH per minute under the conditions described below.

Method:

Reagents

- A. Pyruvate solution : 5.0mM [5.50mg sodium pyruvate (MW=110)/10ml of H₂O] (Should be prepared fresh)
- B. K-Phosphate buffer, pH 7.4 : 1.0M
- C. NADH solution : 1.0mM [7.63mg NADH · 2Na (MW=763)/10ml of H₂O] (Should be prepared fresh)
- D. Enzyme diluent : 0.1M K-phosphate buffer, pH 7.4 contg. 0.1% of BSA

Procedure

1. Prepare the following working solution (10 tests) in a brownish bottle, immediately before use and store on ice.

- | | | |
|--------|----------------------------|-----|
| 3.0ml | Substrate solution | (A) |
| 2.0ml | K-Phosphate buffer, pH 7.4 | (B) |
| 3.0ml | NADH solution | (C) |
| 22.0ml | H ₂ O | (D) |

Concentration in assay mixture		
K-Phosphate buffer	67	mM
Pyruvate	0.49	mM
NADH	0.098	mM
BSA	16.4	μg/mM

2. Pipette 3.0ml of working solution into a cuvette (d=1.0cm) and equilibrate at 25°C for about 5 minutes.
3. Add 0.05ml of the enzyme solution* and mix by gentle inversion.
4. Record the decrease in optical density at 340nm against water for 2 to 3 minutes in a spectrophotometer thermostated at 25°C, and calculate the $\Delta\Delta\text{OD}$ per minute from the initial linear portion of the curve (ΔOD test).

At the same time, measure the blank rate ($\Delta\Delta\text{OD}$ blank) by using the same method as the test except that the enzyme diluent is added instead of the enzyme solution.

* Dilute the enzyme preparation to 0.2–1.0U/ml with ice-cold enzyme diluent (D), immediately before assay.

Calculation

Activity can be calculated by using the following formula :

$$\text{Volume activity (U/ml)} = \frac{\Delta\text{OD/min} (\Delta\text{OD test} - \Delta\text{OD blank}) \times V_t \times df}{6.22 \times 1.0 \times V_s} = \Delta\text{OD/min} \times 9.81 \times df$$

$$\text{Weight activity (U/mg)} = (\text{U/ml}) \times 1/C$$

V_t : Total volume (3.05ml)

V_s : Sample volume (0.05ml)

6.22 : Millimolar extinction coefficient of NADH (cm²/micromole)

1.0 : Light path length (cm)

df : Dilution factor

C : Enzyme concentration in dissolution (c mg/ml)

REFERENCES

- 1) C.A.Loshon, R.B.McComb, L.W.Bond, G.N.Bowers, Jr.W.H.Coleman and R.H.Gwynn; *Clin.Chem.*, **23**, 1576 (1977).
- 2) H.Taguchi, M.Machida, H.Matsuzawa and T.Ohta; *Agric.Biol.Chem.*, **49**(2), 359 (1985).
- 3) F.Gasser, M.Doudoroff, and R.Contopoulos; *J.Gen. Microbiol.* **62**, 241 (1970).

Table 1. Effect of Various Chemicals on D-Lactate dehydrogenase

[The enzyme dissolved in 0.1M K-phosphate buffer, pH 7.4 (20U/ml) was incubated with each chemical at 25°C for 1hr.]

Chemical	Concn.(mM)	Residual activity(%)	Chemical	Concn.(mM)	Residual activity(%)
None	—	100	MIA	2.0	92.4
Metal salt	2.0		NaF	2.0	98.0
MgCl ₂		100	NaN ₃	20	97.2
CaCl ₂		96.3	EDTA	5.0	96.0
Ba(OAc) ₂		95.8	o-Phenanthroline	2.0	98.0
FeCl ₃		94.4	α, α' -Dipyridyl	1.0	97.1
CoCl ₂		97.3	Borate	50	97.5
MnCl ₂		96.9	IAA	2.0	91.6
Cd(OAc) ₂		97.2	NEM	2.0	95.2
NiCl ₂		95.5	Hydroxylamine	2.0	96.3
CuSO ₄		94.3	Triton X-100	0.10%	105.5
Pb(OAc) ₂		96.0	Brij 35	0.10%	104.1
AgNO ₃		71.5	Tween 20	0.10%	106.3
			Span 20	0.10%	98.2
			Na-cholate	0.10%	102.1
			SDS	0.05%	104.4
			DAC	0.05%	47.6

Ac, CH₃CO; MIA, Monoiodoacetate; EDTA, Ethylenediaminetetraacetate; IAA, Iodoacetamide; NEM, N-ethylmaleimide; SDS, Sodium dodecyl sulfate; DAC, Dimethylbenzylalkylammonium chloride.




Fig. 1. Stability (Powder form)
(kept under dry condition, 37°C)




Fig. 2. pH-Activity

[in 57mM buffer solution: pH 4-5, acetate; pH 5-8, K-phosphate; pH 8-9, Tris-HCl]




Fig. 4. pH-Stability

[25°C, 48hr-treatment with 0.1M buffer solution: pH 4-6, dimethylglutaric acid-NaOH; pH 6-8, K-phosphate; pH 8-9, Tris-HCl; pH 9-10, glycine-NaOH. Enzyme concentration: 10U/ml]




Fig. 3. Temperature activity

(in 67mM K-phosphate buffer, pH 7.4)





Fig. 5. Temperature stability

[15min-treatment with 50mM K-phosphate buffer, pH 7.0. Enzyme concentration: 10U/ml]

活性測定法 (Japanese)

1.原理



NADHの消失量を340nmの吸光度の変化で測定する。

2.定義

下記条件下で1分間に1マイクロモルのNADHが酸化される酵素量を1単位(U)とする。

3.試薬

- A. 5.0mM ピルビン酸ナトリウム水溶液(用時調製)
 - B. 1.0M K-リン酸緩衝液, pH7.4
 - C. 1.0mM NADH水溶液(用時調製)
 - D. 酵素溶解液: 0.1M K-リン酸緩衝液, pH 7.4
 - E. 酵素希釈液: 0.1% BSAを含む0.1M K-リン酸緩衝液, pH 7.4
- 酵素溶液: 酵素標品を予め氷冷した酵素溶解液(D)で溶解し,分析直前に酵素希釈液(E)で0.2~1.0U/mlに希釈する。

4.手順

- ① 下記反応混液を使用直前に調製する。
(褐色瓶にて氷冷保存)

3.0ml	基質溶液	(試薬A)
2.0ml	K-リン酸緩衝液	(試薬B)
3.0ml	NADH水溶液	(試薬C)
22.0ml	H ₂ O	
- ② 反応混液3.0mlをキュベット(d=1.0cm)に採り,25℃で約5分間予備加温する。
- ③ 酵素溶液0.05mlを添加し,ゆるやかに混和後,水を対照に25℃に制御された分光光度計で340nmの吸光度変化を2~3分間記録し,その初期直線部分から1分間当たりの吸光度変化を求める(Δ OD test)。
- ④ 盲検は反応混液①3.0mlに酵素溶液の代わりに酵素希釈液(E)0.05mlを加え,上記同様に操作を行って1分間当たりの吸光度変化を求める(Δ ODblank)。

5.計算式

$$\begin{aligned} \text{U/ml} &= \frac{\Delta \text{OD}/\text{min} (\Delta \text{OD test} - \Delta \text{OD blank}) \times 3.05(\text{ml}) \times \text{希釈倍率}}{6.22 \times 1.0 \times 0.05(\text{ml})} \\ &= \Delta \text{OD}/\text{min} \times 9.81 \times \text{希釈倍率} \\ \text{U/mg} &= \text{U/ml} \times 1/\text{C} \\ 6.22 &: \text{NADHのミリモル分子吸光係数} \\ &\quad (\text{cm}^2/\text{micromole}) \\ 1.0 &: \text{光路長(cm)} \\ \text{C} &: \text{溶解時の酵素濃度(c mg/ml)} \end{aligned}$$